

AD 692630

FSTC-HT-23-518-68

U.S. ARMY FOREIGN SCIENCE AND TECHNOLOGY CENTER



DRY, NON-INFECTIOUS, HEMAGGLUTINATING ANTIGENS OF
SIX GROUP A ARBOVIRUSES GROWN IN TISSUE CULTURE

COUNTRY: USSR

TECHNICAL TRANSLATION

This document has been approved
for public release and sale;
its distribution is unlimited.
It may be released to the
Clearinghouse, Department of
Commerce, for sale to the
general public.

TECHNICAL TRANSLATION

FSTC-HT-23- 518-68

Dry, Noninfectious, Hemagglutinating Antigens of Six
Group A Arboviruses Grown in Tissue Culture

P. S. Karaseva, et. al.
Voprosy Virusologii, No. 2, 1967
pages 249-251

Translated for FSTC by ACSI

This translation is an unedited rendition of the original foreign text. Statements or theories advocated or implied are those of the source and do not reflect the position or opinion of the US Army Foreign Science and Technology Center. This translation is published with a minimum of copy editing and graphics preparation in order to expedite the dissemination of information. Requests for additional copies of this document should be addressed to the Defense Documentation Center, Cameron Station, Alexandria, Virginia, ATTN: OSR-2.

UDC 576.858.25.097.34.093.35

DRY, NONINFECTIOUS, HEMAGGLUTINATING ANTIGENS OF
SIX GROUP A ARBOVIRUSES GROWN IN TISSUE CULTURE

P. S. Karaseva, Ye. V. Finogenova, B. F. Semenov,
and I. M. Rodin*

Several authors have shown that hemagglutinins form in growth on tissue cultures of many representatives of group A arboviruses [1-8]. These hemagglutinins have proven highly stable at 37°, which has made possible a reliable method of thermal inactivation of infectious properties without altering the capacity to agglutinate erythrocytes.

In this paper, results of studies performed on various conditions used in drying noninfectious hemagglutinating antigens are presented and experimental features of dry preparations are given.

Materials and Methods

Western (WEE) and Eastern (EEE) equine encephalomyelitis, Sindbis, Semliki forest, Chikungunya, and Middelburg viruses were used.

The WEE, EEE, Sindbis, and Semliki forest viruses were grown in monolayer cultures of chick embryo fibroblasts.. Transplanted cells of kidney epithelium of hamster embryo (VНК-21) were used for accumulation of Chikungunya and Middelburg viruses.

Monolayer cultures were grown in Ru liter separating flasks. Prior to infection, the growth medium was decanted, the monolayer was washed with Hank's solution, after which one ml of virus-containing suspension of mice brain at 1:100 dilution was added to it. The medium No. 199 without serum was used as the maintenance medium.

* Institute of Poliomyelitis and Viral Encephalites of the USSR Academy of Medical Sciences, Moscow

The infected separating flasks were kept in a thermostat for 7 days in the study on Chikungunya and Middelburg viruses, and for 9 days in the study on the WEE, EEE, and Sindbis viruses. Exposure to 32° for 7 days proved optimal for the Semliki forest virus. Under these conditions, as we have shown earlier, maximum accumulation of hemagglutinins and total cessation of infectivity occur [22].

The prepared antigens were dried by the chamber method in the volume after preliminary freezing at temperatures from -60 to 70°. For the first 3 to 4 hours of drying in chamber the temperature was kept at -45 to -35°, then it was gradually raised, and by the 11-12th hour it was raised to 0-1°, and in 24-30 hours -- 20-24°. The drying lasted 28-30 hours at a vacuum of 20-80 microns.

Saccharose and gelatin (final concentration, respectively, 10 and 1 percent), saccharose (final concentration 10 percent), and protamine-sulfate (0.6 mg/ml) were used as the protective mixture.

The hemagglutination reaction [H_h] was established with 0.5 percent suspension goose erythrocytes at room temperature.

Results

Experimental results with WEE, Sindbis, and Middelburg viruses showed that when they are dried without protective mixture the activity of noninfected hemagglutinating antigens is reduced by 16 times (Table 1). When protamine-sulfate was added, the hemagglutinating activity stopped altogether. In the presence of saccharose maintenance of one-half to one-eighth the original titer was noted.

TABLE 1
Activity of Noninfectious Hemagglutinating Antigens
Dried in the Presence of Various Protective
Mixtures

| (A) Antigens | (B) Титр гем-агло- тининов | | | | |
|--------------|-------------------------------|--------------------------|--------------------------|--------------|----------------------------------|
| | (C) Исходный | (D) без напол- нителя | (E) с наполнителем | | |
| (I) | (J) | (K) | (F) протамин- сульфат | (G) сахароза | (H) сахароза с желати- ном |
| | | | | | |
| WEE | 256 | 16 | 0 | 32 | 256 |
| Sindbis | 256 | 16 | 0 | 32 | 256 |
| Middelburg | 128 | 8 | 0 | 64 | 128 |

LEGEND: A - antigen
B - hemagglutination titer
C - original
D - without filler
E - with filler
F - protamine-sulfate
G - saccharose
H - saccharose & gelatin
I - WEE
J - Sindbis
K - Middelburg

TABLE 2

Stability of Liquid and Dry Noninfectious Hemagglutinating
Antigens of Group A Arboviruses at 4 and at 37°

| A Антиген | B Физичес- кое сос- тояние | D Исход- ный | C Активность геммагглютинации после хранения | | | | | | | | | |
|---------------------|-------------------------------------|--------------------|---|------|------|-------|-------|-------------|-------|-------|-------|-------|
| | | | E при 37° | | | | | F при 4° | | | | |
| | | | G Срок хранения (в днях) | | | | | | | | | |
| | | | 15 | 30 | 45 | 60 | 90 | 180 | 1 | 3 | 6 | 12 |
| ЗЭЛ H | N Жидкий | 1:256 | 1:256 | 1:64 | 1:16 | 1:4 | — | — | 1:256 | 1:256 | 1:128 | 1:128 |
| | O Сухой | 1:256 | 1:256 | 1:64 | 1:16 | 1:256 | 1:256 | 1:128 | 1:256 | 1:256 | 1:256 | 1:256 |
| ВЭЛ I | N Жидкий | 1:64 | 1:64 | 1:8 | 1:2 | — | — | — | 1:64 | 1:32 | 1:32 | 1:32 |
| | O Сухой | 1:64 | 1:64 | 1:8 | 1:2 | 1:64 | 1:32 | 1:32 | 1:64 | 1:64 | 1:64 | 1:64 |
| Синдбис J | N Жидкий | 1:128 | 1:128 | 1:32 | 1:8 | 0 | — | — | 1:128 | 1:128 | 1:128 | 1:128 |
| | O Сухой | 1:128 | 1:128 | 1:32 | 1:8 | 1:128 | 1:128 | 1:64 | 1:128 | 1:128 | 1:128 | 1:128 |
| Леса Семли- ки K | N Жидкий | 1:64 | 1:64 | 1:8 | 1:2 | — | — | — | 1:64 | 1:64 | 1:16 | 1:16 |
| | O Сухой | 1:64 | 1:64 | 1:8 | 1:2 | 1:64 | 1:32 | 1:32 | 1:64 | 1:64 | 1:32 | 1:32 |
| Чикунгунья L | N Жидкий | 1:64 | 1:64 | 1:8 | 1:2 | 0 | — | — | 1:64 | 1:64 | 1:64 | 1:64 |
| | O Сухой | 1:64 | 1:64 | 1:8 | 1:2 | 1:64 | 1:64 | 1:32 | 1:64 | 1:64 | 1:64 | 1:64 |
| Мидделбург M | N Жидкий | 1:128 | 1:128 | 1:16 | 1:2 | 0 | — | — | 1:128 | 1:128 | 1:128 | 1:128 |
| | O Сухой | 1:128 | 1:128 | 1:16 | 1:2 | 1:128 | 1:128 | 1:64 | 1:128 | 1:128 | 1:64 | 1:64 |

LEGEND: A - antigen
B - physical condition
C - hemagglutination titer after storage
D - original
E - at 37°
F - at 4°
G - length of storage (in days)
H - WEE
I - EEE
J - Sindbis
K - Semliki forest
L - Chikungunya
M - Middelburg
N - liquid
O - dry

The mixture of saccharose and gelatin gave the most pronounced protective effect. It made it possible to get dry antigens without variation in the initial activity in the HR.

Several series of antigens of each viruses were obtained with this protective mixture.

Table 2 lists results in the determination of the hemagglutinating activity of liquid and dry antigens during storage and at 4 and 37°. It was found that liquid antigens of most of the viruses studied are highly stable at 4°. Titers of the antigens -- WEE, EEE, Sindbis, and Chikungunya -- remained practically unchanged for the course of the year (the

course of the observation period).

The Middelburg antigen retained all its activity for 9 months (observation period). Hemagglutinins of Semliki forest virus showed less stability. After 6 months of storage, a fourfold reduction was noted. At 37° all antigens tested quickly lose the ability to agglutinate erythrocytes. In 15 days, the antigen of Semliki forest virus becomes inactive, and in 30 days -- the antigens of Sindbis, Chikungunya, and Middelburg become inactive. In a month, the titer of hemagglutinins of WEE virus was reduced by 64 times.

Dry antigens show much stability not only at 4, but also at 37°. Under thermostat conditions their activity remained unchanged for 3 months in experiments with all the viruses. After 6 months, using the models of the viruses WEE, Sindbis, Middelburg, and Chikungunya, only a twofold titer reduction was observed.

Conclusions

1. An optimal regime for lyophilization of noninfectious hemagglutinating antigens of group A arboviruses grown on tissue cultures has been developed.

2. The high stability of dry antigens at 37° for a three-month period has been demonstrated.

BIBLIOGRAPHY

1. Gnydamovich, S. Ya., Obukhova, V. R. et al, in: Materialy 17-y Sessii In-ta Virusologii im. D. I. Ivanovskogo (Materials of the 17th Session of the Institute of Virology imeni D. I. Ivanovskiy), Moscow, 1964, Part 2, page 45.

2. Karasova, P. S., Semenov, B. F., et al. Aktual'nyye Problemy Virusnykh Infektsiy (Urgent Problems in Viral Infections), Moscow, page 129, 1965.

3. Rzepova, A. I., Salagova, T. A., and Neustroyev, V. D., *ibid.*, page 163.

4. Buckley, S. M. and Srihongse, A. S., Proc. Soc. Exp. Biol. (N.Y.), Vol. 113, page 284, 1963.

5. Likar, M., Buckley, S. M., and Clarke, D. H., Virology, Vol. 18, page 640, 1962.

6. Pattyn, S. R. and De Vleeschauwer, L., Ann. Soc. Belge Med. Trop., Vol. 44, page 525, 1964.
7. Salminen, A., Ann. Med. Exp. Fenn., Vol. 40, page 174, 1962.
8. Idem, Acta Path. Microbiol. Scand., Suppl. 154, page 343, 1962.

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

| | | | |
|---|-----------------------------|---|--|
| 1. ORIGINATING ACTIVITY (Corporate author) Foreign Science and Technology Center US Army Materiel Command Department of the Army | | 2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED | |
| 3. REPORT TITLE DRY, NONINFECTIOUS, HEMAGGLUTINATING ANTIGENS OF SIX GROUP A ARBOVIRUSES GROWN IN TISSUE CULTURE | | 2b. GROUP | |
| 4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Translation | | | |
| 5. AUTHOR(S) (First name, middle initial, last name) P. S. Karaseva, et al. | | | |
| 6. REPORT DATE | 7a. TOTAL NO. OF PAGES 5 | 7b. NO. OF REFS N/A | |
| 8a. CONTRACT OR GRANT NO. | | 8b. ORIGINATOR'S REPORT NUMBER(S) FSTC-HT-23-518-68 | |
| a. PROJECT NO. | | 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) | |
| c. 02TR100 | | ACSI Control Number J-6365 | |
| d. SNUFD-PR-I | | | |
| 10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited. | | | |
| 11. SUPPLEMENTARY NOTES | | 12. SPONSORING MILITARY ACTIVITY US Army Foreign Science and Technology Center | |
| 13. ABSTRACT In this paper, results of studies performed on various conditions used in drying noninfectious hemagglutinating antigens are presented and experimental features of dry preparations are given. | | | |

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

UNCLASSIFIED
Security Classification

LINK A

LINK

LINK C

ROLE**WOL**

www

ROLE

UNCLASSIFIED